



## Key Targets

**1** Reduce pre-weaning mortality to below 9.3%

(Agrosoft data 2015 – top 10% UK indoor breeding herds)



**2** Increase pigs weaned per litter to at least 12.5

(Agrosoft data 2015 – top 10% UK indoor breeding herds)




**3** Improve weaning weight, aim for 8kg at 28 days



# Newborn management (indoors)

**Newborn piglets are poorly equipped to keep warm immediately after birth and, as the energy reserves to produce body heat (glycogen stored in the liver) are limited, the combination of factors below means that piglets often become chilled.**

Piglets are also born with no immunity and can only get this from colostrum soon after birth. A minimum of **100ml** of colostrum per kilogram of birthweight within the first **16** hours is crucial to provide the energy, nutrients and antibodies needed for survival.



### HOW CAN YOU HELP THE NEWBORN PIGLET AND ENSURE A GOOD START IN LIFE?

#### Monitor farrowing

Being present at farrowing is a critical part of good farrowing house management. If not possible on your unit, ask yourself why not? Consider altering routines or rotas to provide cover at this crucial time or discuss with your vet the practicalities of using products to promote farrowing during the daytime, when you can be present.

#### Colostrum

- Colostrum is the 'first milk' and an essential source of energy, nutrients and immunity for the piglet
- Colostrum is critical for development of the piglets' own immune system and optimum lifetime performance
- Maximise colostrum intake in the first six hours after birth, achieving at least **100ml** intake per kilogram of birthweight by **16** hours post farrowing. For example, a **1.45kg** piglet would need a minimum of **145ml** colostrum
- Beyond **24** hours is too late, as the piglets' intestines can no longer absorb the large antibodies found in colostrum
- Colostrum is only available in quantity for about **12** hours and after **20** hours the sow will be producing milk and not colostrum.

### FOSTERING MANAGEMENT

#### Cross fostering

Make sure all staff are aware of your fostering policy. Sometimes, practices change for the better but can revert back if your cover staff are not aware of the changes. Wherever possible, leave piglets with their own mothers to avoid disruption of the litter suckling patterns; even big piglets looking secure and strong will experience growth checks if fostered, especially if moved around frequently.

#### Even up numbers of piglets per litter

Matching the number of piglets to the number of functioning teats allows each piglet to have easy access to the colostrum produced after farrowing. This improves piglet survival rates and increases the chance of piglets achieving their potential growth rates.

#### Helping low birthweight piglets

It is a great disadvantage for small piglets to have to compete with larger littermates. Litters of small piglets should be created from all the 'smalls' born in a given farrowing day. Foster litters of small piglets should be put with low-parity sows; the teat size of a low-parity sow will match the small mouths of the small piglets.



### RULES OF FOSTERING

- Foster within the first **24** hours
- Ensure foster piglets get early colostrum, referably from their own mother, or, if not, from a very recently farrowed foster mother
- Give the smaller piglets warm colostrum (sow, artificial or pasteurised bovine) or glucose
- Foster only once, it can be detrimental to keep moving litters around
- Remember, small piglets find suckling from large teats difficult

- Never have more piglets than functioning teats on sows
- Foster within a room (batch of piglets), wherever possible, don't move health problems to other groups.

### OBSERVE PIGLET BEHAVIOUR

- Look out for fading piglets and take remedial action by fostering and providing electrolytes or additional milk
- Check for signs of ill health and treat according to unit veterinary protocol
- Ensure piglets are lying either on or under the heat source and not piling together, indicating chilling.

**Remember:**  
Colostrum deprivation will reduce lifelong survivability and lifelong performance, see Table 1.

**Table 1: Ensuring colostrum intake**

ACTIONS	WHEN TO USE IT	HOW	TIME TAKEN
Split suckling	Litter size is large. Considerable variation in piglet size.	Split the litter into two groups. Initially, enclose the group of heavier, stronger piglets within the creep area, behind a board or in a box, to reduce competition. Allow the smaller, less viable piglets to suckle and then swap the groups over after <b>90 minutes</b> to allow for two sucklings at <b>40-minute</b> intervals. Both groups of piglets should be able to access a warm environment.	On average, <b>five minutes</b> for each litter requiring extra help. It is an easy task and can be combined with litter work.
Assisted suckling	High numbers of small, low viability piglets have been born. Considerable variation in piglet size.	Supervise and assist the vulnerable piglets to gain access to teats and suckle. Ensure they are able to suckle unhindered, then mark each piglet once seen to suckle. Consider stomach tubing vulnerable piglets.	On average, <b>10–15 minutes</b> for each litter requiring extra help. This requires considerable patience but can be combined with split suckling.
Hand feeding colostrum	High numbers of small, low viability piglets have been born. When creating litters of small piglets.	Milk sows that have farrowed within the last <b>five hours</b> . Syringe feed the piglets with the colostrum when establishing the new litter until the suckling pattern is established (the colostrum can be fresh or you can store frozen colostrum and defrost naturally as required).	On average, allow at least <b>15 minutes</b> for milking and feeding the piglets that require the extra help. It can be time-consuming but colostrum is essential for piglet survival.



Split suckling

### CREEP TRAINING

Piglets have different thermal requirements from sows, the creep area provides the required environment for the piglets and training them to use this area will provide them with a good start in life. The creep has a secondary advantage of being a safe area where piglets can lie away from the sow and reduce the risk of being laid on.

- Piglets learn the behaviour of using creeps within the first **40–48** hours of life
- Enclose the piglets within a closed creep as soon as they have finished suckling
- This can be combined with split suckling, saving time by carrying out the two activities at the same time
- Keep piglets in the creep area for approximately **one** hour.

© Agriculture and Horticulture Development Board 2017. No part of this publication may be reproduced in any material form (including by photocopy or storage in any medium by electronic means) or any copy or adaptation stored, published or distributed (by physical, electronic or other means) without the prior permission in writing of the Agriculture and Horticulture Development Board, other than by reproduction in an unmodified form for the sole purpose of use as an information resource when the Agriculture and Horticulture Development Board is clearly acknowledged as the source, or in accordance with the provisions of the Copyright, Designs and Patents Act 1988. All rights reserved.

While the Agriculture and Horticulture Development Board seeks to ensure that the information contained within this document is accurate at the time of printing, no warranty is given in respect thereof and, to the maximum extent permitted by law, the Agriculture and Horticulture Development Board accepts no liability for loss, damage or injury howsoever caused (including that caused by negligence) or suffered directly or indirectly in relation to information and opinions contained in or omitted from this document.

AHDB Pork is part of the Agriculture and Horticulture Development Board.